

## Family 127. CONNARACEAE

### Revision:

Leenhouts, PW. 1958. Connaraceae. Flora Malesiana series 1, 5: 495-541.

### 1. Agelaea Solander ex Planchon

**Agelaea borneensis** (Hook.f.) Merr., PJS 4 c (1909) Bot. 127; --Leenh., Fl. Males. ser. 1, 5 (1958) 504, Fig. 2i.

Agelaea everettii Merr., PJS 4 c (1909) Bot. 127; EPFP 2 (1923) 239;

Burma?, Sumatra, Peninsular Malaysia, Borneo, Philippines, W & C Java.

Primary and secondary rainforests,

**Agelaea trinervis** (Llanos) Merr., Sp. Blancoanae (1918) 164; EPFP 2 (1923) 239; --Leenh., Fl. Males. ser. 1, 5 (1958) Hainan, Thailand, Laos, Cambodia, C & S Vietnam, Sumatra, Peninsular Malaysia, Borneo, Philippines, W Java, Moluccas (Sula Is; Seram).

### 2. Cnestis Jussieu

### Revision:

Andreas, Ch H & N Prop. 1954. *Florae Malesianae precursors VIII. The genus Cnestis (Connaraceae) in Indo-Malaysia*. Blumea 7: 602-616.

**Cnestis palala** (Lour.) Merr., J. Straits Branch Roy. As. Soc. no. 85 (1922) 201; EPFP 2 (1923) 240.  
**ssp. diffusa** (Blanco) Andreas, Blumea 7 (1954) 613; --Leenh., Fl. Males. ser. 1, 5 (1958) 499, Fig. 1e-j.

Sumatra (E coast), Lingga Archipelago, Peninsular Malaysia, Borneo, Philippines.

LUZON, LUBANG, SEMIRARA, BURIAS, CEBU. Rainforests and thickets, along streams, up to 200m. Fl. (Oct.-) Jan.-March (-April), fr. May-July.

### 3. Connarus Linnaeus

**Connarus culionensis** Merr., PJS 4 c (1909) Bot. 120; EPFP 2 (1923) 237; --Leenh., Fl. Males. ser. 1, 5 (1958) 531. --Type:

Connarus erianthus Elmer, LPB 5 (1913) 1762; --Merr., EPFP 2 (1923) 237. --Type:

#### **var. culionensis**

S Vietnam, Borneo (Sarawak; Sabah), Banguey, Philippines.

CALAMIANES, PALAWAN, BALABAC. Dry open slopes, dry thickets and forests at low altitude.

**var. stellatus** (Merr.) Leenh., Fl. Males. ser. 1, 5 (1958) 533; --Connarus stellatus Merr., PJS 4 c (1909) Bot. 119; EPFP 2 (1923) 238. --Type:

Distribution as the preceding.

**Connarus grandis** Jack, Mal. Misc. 2 no. 7 (1822) 40; --Leenh., Fl. Males. ser. 1, 5 (1958) 529, Fig. 13g.

Connarus trifoliatus (Turcz.) Rolfe, J. Bot. 23 (1885) 212, (as 'trifoliolatus'); --Merr., EPFP 2 (1923) 238.

--Type:

Sumatra, Simalur, Banka, Peninsular Malaysia, Borneo, Philippines, possibly Talaud Isls, W Java.

**Connarus monocarpus** L., Sp. Pl. (1753) 675;

**ssp. malayensis** Leenh., Fl. Males. ser. 1, 5 (1958) 538.

Connarus hallieri Merr., PJS 4 c (1909) Bot. 122; EPFP 2 (1923) 237. –Type:

Connarus palawanensis Elmer, LPB 5 (1913) 1763; --Merr., EPFP 2 (1923) 238. –Type:

Nicobar Isls., throughout Malesia (except the Moluccas and New Guinea).

LUZON: Quezon, Laguna, Camarines, Sorsogon, PALAWAN, SIBUYAN, LEYTE, SAMAR, BASILAN. Lowland and medium altitude forests.

**Connarus semidecandrus** Jack, Mal. Misc. 2 no. 7 (1822) 39; --Leenh., Fl. Males. ser. 1, 5 (1958) 534, Fig. 11d, g-h.

Connarus balsahanensis Elmer, LPB 5 (1913) 1764; --Merr., EPFP 2 (1923) 237.

Connarus neurocalyx Planch., Linnaea 23 (1850) 428; --Merr., Fl. Manila (1912) 220; EPFP 2 (1923) 238.

Andaman Isls, Burma, Thailand, S Indochina, throughout Malesia (except E half of Java), also in Micronesia (Palau) and Melanesia (Solomon Isls), also Truk.

LUZON (Isabela) southward to PALAWAN, PANAY and NEGROS. Common in lowland forests and thickets.

**Connarus subfoveolatus** Merr., PJS 13 c (1918) Bot. 15; EPFP 2 (1923) 238; --Leenh., Fl. Males. ser. 1, 5 (1958) 541. –Type: BS 28240 Fenix (NY\*, US\*, iso). Luzon: Apayao prov., May 1917.

LUZON: Apayao. Damp forests at medium altitude. Fl. May.

**Connarus subinaequifolius** Elmer, LPB 1 (1908) 297; --Merr., EPFP 2 (1923) 238; --Leenh., Fl. Males. ser. 1, 5 (1958)

**var. subinaequifolius**

Connarus bracteatus Merr., PJS 4 c (1909) Bot. 120; EPFP 2 (1923) 237. –Type:

LUZON, DINAGAT. Forests up to 850m. Fl. Apr-May, fr. Mar-May.

**var. sericeus** Leenh., Fl. Males. ser. 1, 5 (1958) 530. –Type:

Connarus castaneus Merr., PJS 14 (1919) 403; EPFP 2 (1923) 237. –Type:

BATAN, LUZON, POLILLO. Low altitude forests, Fr. Nov-Dec.

**Connarus whitfordii** Merr., PJS 4 c (1909) Bot. 123; EPFP 2 (1923) 238; --Leenh., Fl. Males. ser. 1, 5 (1958) 536, Fig. 14. –Type: FB 9158 Whitford & Hutchinson

Connarus caudatus Merr., PJS 17 (1920) 261; EPFP 2 (1923) 237. –Type: BS 35059 Ramos & Pascasio

Connarus mindanaensis Merr., PJS 4 c (1909) Bot. 122; EPFP 2 (1923) 237. –Type:

Connarus oliganthus Elmer, LPB 10 (1939) 3718. –Type: Elmer 16750

LUZON: Sorsogon (Mt Bulusan), POLILLO, BILIRAN, LEYTE, SAMAR, MINDANAO:

Zamboanga, BUCAS GRANDE. In and along primary and secondary forests, up to c. 750m. Fl. Jan-Apr (Jul, Sept), fr. Oct-Jan (Apr, Jun).

**Connarus winkleri** Schellenb., Bot. Jahrb. 59 (1924) Beibl. no. 131, p. 138

**ssp. philippinensis** Leenh., Blumea Suppl. 4 (1958) 106; Fl. Males. ser. 1, 5 (1958) 540. –Type: BASILAN.

#### 4. **Ellianthus** JD Hooker

**Ellianthus tomentosus** Kurz, J. As. Soc. Beng. 41, 2 (1872) 305;

**ssp. tomentosus var. luzoniensis** (Vidal) Leenh., Fl. Males. ser. 1, 5 (1958) 521, Fig. 10b-e; --

Soejarto et al. (1995) Prelim. Checklist Fl. Pl. Palawan; --Ellianthus luzoniensis Vidal, Rev. Pl.

Vasc. Filip. (1886) 104; --Merr., EPFP 2 (1923) 241. –Type:

*Ellipanthus longifolius* Merr., PJS 17 (1920) 262; EPFP 2 (1923) 241. –Type:  
*Ellipanthus mindanaensis* Merr., PJS 4 c (1909) Bot. 124; EPFP 2 (1923) 241. –Type:  
*Ellipanthus urdanetensis* (Elmer) Merr., EPFP 2 (1923) 241. –Type:  
Borneo (Sarawak), Philippines, C Sulawesi.  
LUZON: Cagayan, Bataan, Bulacan, Quezon, Camarines, Sorsogon, PALAWAN,  
MINDANAO: Zamboanga, Davao, Agusan, DINAGAT. Rainforests, from lowlands to c.  
1100m.

## 5. *Rourea* Aublet

***Rourea minor* (Gaertn.) Alston**, Handb. Fl. Ceylon 6 (Suppl.) (1931) 67; --Leenh., Fl. Males. ser. 1, 5  
(1958) 514, Fig. 8.

*Rourea erecta* (Blanco) Merr., PJS 4 c (1909) Bot. 125; Fl. Manila (1912) 220; EPFP 2 (1923) 239. –Type:  
*Rourea imbricata* Elmer, LPB 7 (1915) 2597; --Merr., EPFP 2 (1923) 240. –Type: Elmer 14309 (NA\*, iso; but  
replicates also distributed as '13951' in L\*, MO\*, NY\*, US\*)  
*Rourea luzoniensis* Merr., PJS 14 (1919) 404; EPFP 2 (1923) 240. –Type:  
*Rourea unifoliolata* Merr., PJS 8 c (1913) Bot. 372; EPFP 2 (1923) 240. –Type:  
*Rourea volubilis* (*non* *Cnestis volubilis* Blanco) Merr., Govt. Lab. Publ. (Philipp.) 27 (1905) 36; EPFP 2  
(1923) 240.

Sri Lanka, the Andaman and Nicobar Isls, SW Deccan, continental SE. Asia from  
Bangladesh and Assam to S China (Hongkong), Hainan and Kotosyo Isl. near Taiwan.  
Across Malesia (except the Lesser Sunda Islands east of Bali). to NE Queensland, New  
Caledonia, Vanuatu, Fiji and Samoa Isls, Solomon Isls., Tonga, Niue I.  
Widespread throughout the Philippines. N to C LUZON: Cagayan to Laguna, Camarines,  
LUBANG, MINDORO, PALAWAN, CUYO, BANTAYAN, PANAY, LEYTE, MINDANAO:  
Agusan. In primary and secondary forests, dry thickets, also more open places and  
along riverbanks. Usually a large liana, rarely shrub or treelet, from the lowlands to c.  
950m.