

Family 7. GNETACEAE

Revision:

Markgraf, F. 1951. *Gnetaceae*. Flora Malesiana series 1,4: 338-347; idem. 1, 6 (1972) 944-949.

1. *Gnetum* Linnaeus

Gnetum arboreum Foxw., PJS 6 c (1911) Bot. 174, t. 32; --Merr., EPFP 1 (1922) 7; --Markgraf, Fl. Males. ser. 1, 4 (1951) LUZON.

Gnetum gnemon L., Mant. 1 (1867) 125; --Merr., EPFP 1 (1922) 7; --Markgraf, Fl. Males. ser. 1, 4 (1951) 340.

var. gnemon

Fiji and Solomon Isls to Malesia: from Sumba and Sulawesi through the Philippines to New Guinea, the Malay Peninsula, and possibly elsewhere, often planted and frequently naturalized in secondary forests, even in W Malesia.

Gnetum gnemonoides Brongn. in Duperry Voy. Bot. (1829) 12; --Merr., EPFP 1 (1922) 7; --Markgraf, Bull. Jard. Bot. Buitenz. 3, 10 (1930) 480, t. 13; Fl. Males. ser. 1, 4 (1951) 344, Fig. 5. New Hanover, Peninsular Malaysia (Taiping), Billiton, Borneo, Philippines, Sulawesi, Moluccas (Aru Isls), New Guinea. Absent from the outer arch of islands, viz. Sumatra, Java, Lesser Sunda Isls.

Gnetum latifolium Blume, Tijd. Nat. Geschied. & Phys. 1 (1834) 160; --Markgraf, Bull. Jard. Bot. Buitenz. 3, 10 (1930) 458; Fl. Males. ser. 1, 4 (1951) 342, Fig. 1g-h. *Gnetum indicum* (Lour.) Merr., Interp. Herb. Amboin. (1917) 77; EPFP 1 (1922) 7, *in part*.

var. latifolium forma latifolium

Philippines, New Guinea.

var. longipes Markgraf, Bull. Jard. Bot. Buitenz. 3, 10 (1930) 459, t. 7, Fig. 7; Fl. Males. ser. 1, 4 (1951) 342.

forma longipes

Indochina, Philippines, New Guinea. LUZON.

var. laxifrutescens (Elmer) Markgraf, Bot. Jahrb. 60 (1925) 148; Bull. Jard. Bot. Buitenz. 3, 10 (1930) 463, t. 7; Fl. Males. ser. 1, 4 (1951) 342, Fig. 1g. Philippines and Moluccas (Kei & Tanimber Is).

var. minus (Foxw.) Markgraf, Bull. Jard. Bot. Buitenz. 3, 10 (1930) 463; Fl. Males. ser. 1, 4 (1951) 343.

Gnetum minus Foxw., PJS 6 c (1911) Bot. 176, t. 33; --Merr., EPFP 1 (1922) 8. Indochina, Borneo (Sabah, Mt Kinabalu), Philippines, SE Sulawesi.