

Family 182. OCHNACEAE

Revision:

Kanis, A. 1971. *Ochnaceae*. Flora Malesiana series 1, 7: 97-119.

1. *Brackenridgea* A Gray

Brackenridgea fascicularis (Blanco) Fern.-Villar, Nov. App. (1880) 40; --Kanis, *Blumea* 16 (1968) 43; Fl. Males. ser. 1, 7 (1971) 104; --*Ochna fascicularis* Blanco, Fl. Filip. ed. 2 (1845) 245; *ibid.* ed. 3, 2 (1878) 92; --Vidal, *Sinopsis Atlas* (1883) 19, t. 27A; Rev. Pl. Vasc. Filip. (1886) 79; --Merr., EPFP 3 (1923) 68.

ssp. *fascicularis*

LUZON: Bulacan, Rizal, Laguna, Quezon, Camarines, SIBUYAN, MINDANAO: Surigao. Lowland and medium altitude primary forests.

ssp. *mindanaensis* (Merr.) Kanis, *Blumea* 16 (1968) 44; Fl. Males. ser. 1, 7 (1971) 104; --*Ouratea mindanaensis* Merr., PJS 17 (1920) 287; EPFP 3 (1923) 68. --Type: BS 34479 Ramos & Pascasio. MINDANAO: Surigao del Norte. Lowland thickets or forests.

Brackenridgea palustris Bartell., *Malpighia* 15 (1901) 165, t. 10.

ssp. *foxworthyi* (Elmer) Kanis, *Blumea* 16 (1968) 48; Fl. Males. ser. 1, 7 (1971) 102; --*Ochna foxworthyi* Elmer, LPB 5 (1913) 1823; --Merr., EPFP 3 (1923) 68. --Type: PALAWAN. Lowlands, along river in forest or rocky hillside near riverbank, 0-150m.

Gomphia Schreber

Gomphia serrata (Gaertn.) Kanis, *Taxon* 16 (1967) 422; *Blumea* 16 (1968) 33; Fl. Males. ser. 1, 7 (1971) 105, Fig. 3; --*Meesia serrata* Gaertn., *Fruct.* 1 (1788) 344, t. 70, Fig. 6.

Ouratea angustifolia (Vahl) Baill. ex Laness., *Pl. Util. Col. Fr.* (1886) 607; --Merr., EPFP 3 (1923) 68.

SW Peninsular India, Sri Lanka, E Thailand, Indochina, Hainan, Sumatra, Banka, Peninsular Malaysia, Karimundjawa Isls, Borneo, Philippines, Sulawesi.

LUZON: Ilocos Norte, Cagayan, Zambales, Quezon, Camarines, MINDORO, CULION, BUSUANGA, PALAWAN, BALABAC, MASBATE, ROMBLON, PANAY, GUIMARAS, NEGROS, MINDANAO. Lowland primary forests, often common.

Neckia Korthals

Neckia serrata Korth., *Ned. Kruidk. Arch.* 1 (1848) 358; --Kanis, *Blumea* 16 (1968) 69; Fl. Males. ser. 1, 7 (1971) 110, Fig. 5.

Neckia philippinensis Merr. & Quisumb., PJS 82 (1954) 329, t. 2. --Type:

Sumatra, Mentawai Isls, Riouw & Lingga Isls, SE Peninsular Malaysia, Borneo & CE Philippines.

SAMAR.

Ouratea Aublet = *Brackenridgea*

Schuermansia Blume

Schuermansia elegans Blume, *Ann. Mus. Bot. Lugd.-Bat.* 1 (1850) 177, t. 32; --Kanis, *Nova Guinea, Bot.* 6 (1961) 64, Fig. 1a; *Blumea* 16 (1968) 75; Fl. Males. ser. 1, 7 (1971) 115.

Borneo, SE Philippines, Sulawesi, Moluccas, and New Guinea.
MINDANAO: Agusan, a single collection of uncertain identity.

Schuermansia vidalii (Fern.-Villar) Merr., PJS 11 c (1916) Bot. 19; --Merr., EPFP 3 (1923) 69; --
Kanis, Nova Guinea, Bot. 6 (1961) 69, Fig. 1c; Blumea 16 (1968) 76; Fl. Males. ser. 1, 7 (1971)
116; --Calophyllum vidalii Fern.-Villar in Ceron, Cat. Pl. Herb. Manila (1892) 229, plate.

Schuermansia parvifolia Merr., PJS 11 c (1916) Bot. 19. --Type:

LUZON: Aurora, Camarines Sur, Sorsogon. In (mossy) forest, altitude 500 to well over
1000m, possibly also at sea-level.

Cultivated:

Ochna kirkii Oliv., Fl. Trop. Afr. 1 (1868) 317.