

Family 224. ONAGRACEAE

Revision:

Raven, PH. 1977. *Onagraceae*. Flora Malesiana series 1, 8: 98-113.

1. *Epilobium* Linnaeus

Revision:

Raven, PH. 1967. The genus *Epilobium* in Malasia (Onagraceae). *Blumea* 15: 269-282.

***Epilobium brevifolium* D Don**, Prodr. Fl. Nepal. (1825) 222

ssp. *trichoneurum* (Haussknecht) Raven, Bull. Br. Mus. Nat. Hist. Bot. 2 (1962) 362; *Blumea* 15 (1967) 271; Fl. Males. ser. 1, 8 (1977) 109; --*Epilobium trichoneurum* Haussknecht, Oest. Bot. Z. 29 (1879) 54.

Epilobium philippinense C Robinson, PJS 5 c (1910) Bot. 369; --Merr., EPFP 3 (1923) 220, excl. syn. E. *sarmentaceum* Hausskn. --Type: Merrill 4484 (K, iso). Luzon: Mountain Province, Mt Data.

SE Tibet to Assam, Myanmar, and W China, N Philippines. LUZON: Mountain Province (Mt Data), Benguet (Mankayan to Baguio; Ambuklao to Baguio; Pauai; Mt Pulag; Mt Sto Tomas; Baguio). Wet open slopes in the pine region, along streams and by springs, altitude 1400-2100m. Fl. Jul, Oct.

***Epilobium platystigmatosum* C Robinson**, PJS 3 c (1908) Bot. 210; --Merr., EPFP 3 (1923) 221; --Raven, *Blumea* 15 (1967) 272; Fl. Males. ser. 1, 8 (1977) 110. --Type: Merrill 4744

Japan, China (Hubei; Jiangsu), Taiwan, N Philippines. LUZON: Ifugao, Mountain Province, Benguet. Chiefly along small streams and about cliffs, altitude 1200-2400m. Fl. Apr-Jun.

Jussiaea Linnaeus =Ludwigia

2. *Ludwigia* Linnaeus

Revision:

Raven, PH. 1963. *Reinwardtia* 6: 327-

***Ludwigia adscendens* (L.) Hara**, J. Jap. Bot. 28 (1953) 290; --Raven, *Reinwardtia* 6 (1963) 387, maps 31, 33; Fl. Males. ser. 1, 8 (1977) 104, Fig. 5; 6 (map); --*Jussiaea adscendens* L., Mantissa 1 (1767) 69.

Jussiaea repens L., Sp. Pl. (1753) 388; --Merr., EPFP 3 (1923) 219; --Backer & Bakh.f., Fl. Java 1 (1963) 260.

Sri Lanka, India, Bangladesh, S China, Hainan, Myanmar, Thailand, Indochina, across Malesia to Guinea, N Australia (W Arnhem Land). Freshwater pools, ditches, swamps, fallow and wet paddies, very common, from lowlands up to **xxxx**

***Ludwigia hyssopifolia* (G Don) Exell**, Garcia de Orta 5 (1957) 471; --Raven, *Reinwardtia* 6 (1963) 387, maps 30; Fl. Males. ser. 1, 8 (1977) 104, Fig. 4; --*Jussiaea hyssopifolia* G Don, Gen. Syst. 2 (1832) 693.

Jussiaea linifolia Vahl, Eclog. Am. 2 (1798) 32; --Merr., Fl. Manila (1912) 356; EPFP 3 (1923) 219; --Backer & Bakh.f., Fl. Java 1 (1963) 260.

Tropical Africa, continental SE Asia (Sri Lanka to Hainan), throughout Malesia to Micronesia and N Australia (Cape York Peninsula and Arnhem Land). Pools, along rice paddies, canals and shallow ditches, river edges, waterlogged wastelands, etc. Lowlands to c. **xxxx** m.

Ludwigia octovalvis (Jacq.) Raven, Kew Bull. 15 (1962) 476; Reinwardtia 6 (1963) 356, maps 19-20, incl. ssp. *brevisepala* (Brenan) Raven & ssp. *sessiliflora* (Mich.) Raven; Fl. Males. ser. 1, 8 (1977) 101, Fig. 1; --*Jussiaea octovalvis* (Jacq.) Sw., Obs. Bot. (1791) 142; --*Oenothera octovalvis* Jacq., En. Syst. Pl. (1760) 19.

Ludwigia villosa Lam., Encycl. 3 (1789) 331; --Merr., EPFP 3 (1923) 220.

Jussiaea erecta L., Sp. Pl. (1753) 388; --Merr., EPFP 3 (1923) 219.

Jussiaea suffruticosa Miq., Fl. Ind. Bat. 1¹ (1856) 628; --Merr., Fl. Manila (1912) 357.

Pantropic, between c. 32°N and 30°S. Waterlogged sites, pools, rice fields, along water-courses, lake shores, also gravelly riverbeds, etc.

Ludwigia perennis L., Sp. Pl. (1753) 119; --Raven, Fl. Males. ser. 1, 8 (1977) 103, Fig. 2 (map); --Raven, Reinwardtia 6 (1963) 367, map 21.

Ludwigia parviflora Roxb., Hort. Beng. (1814) 11, *nom. nud.*; Fl. Ind. 1 (1820) 440; --Merr., Fl. Manila (1912) 355; EPFP 3 (1923) 220.

Tropical Africa, Madagascar, Sri Lanka to S China, Hongkong, and Hainan, throughout Malesia (except Borneo and Moluccas) to tropical Australia and New Caledonia. Sunny waterlogged conditions, along fallow rice-fields, watercourses, riverbeds, on heavy clay, etc. from lowlands to c. 500m. Fl. throughout the year.

Ludwigia prostrata Roxb., Hort. Beng. (1814) 11, *nom. nud.*; Fl. Ind. 1 (1820) 441; --Merr., Fl. Manila (1912) 355; EPFP 3 (1923) 220; --Raven, Reinwardtia 6 (1963) 374, map 23; Fl. Males. ser. 1, 8 (1977) 103, Fig. 3 (map);

Sri Lanka and S Deccan to N India, Assam, Andaman Isls, S China, Peninsular Malaysia, Java, Lesser Sunda Isls (Timor), Borneo, and the Philippines. LUZON, PALAWAN, NEGROS, MINDANAO. Lowlands to c. 800m, along rice paddies, ditches, river banks, swampy sites, rather rare.

Cultivated:

Fuchsia magellanica Lam. var. **gracilis** (Nich.) Bailey; Backer & Bakh.f., Fl. Java 1 (1963) 264.

Native of temperate South America.